arXiv:1207.3693v1 [math.DS] 16 Jul 2012

STATISTICAL STABILITY FOR MULTI-SUBSTITUTION TILING SPACES

RUI PACHECO AND HELDER VILARINHO

ABSTRACT. Given a finite set $\{S_1 \ldots, S_k\}$ of substitution maps acting on a certain finite number (up to translations) of tiles in $\mathbb{R}^d$, we consider the multi-substitution tiling space associated to each sequence $\bar{a} \in \{1, \ldots, k\}^{\mathbb{N}}$. The action by translations on such spaces gives rise to uniquely ergodic dynamical systems. In this paper we investigate the rate of convergence for ergodic limits of patches frequencies and prove that these limits vary continuously with $\bar{a}$.

1. INTRODUCTION

Roughly speaking, a *tiling* of $\mathbb{R}^d$ is an arrangement of tiles that covers $\mathbb{R}^d$ without overlapping. An important class of tilings is that of *self-similar tilings*. In order to construct a self-similar tiling x, one starts with a finite number (up to translation) of tiles and a *substitution map* S that determines how to inflate and subdivide these tiles into certain configurations of the same tiles. Many examples can be found in [3, 8]. The *substitution tiling space* X_S is the closure of all the translations of x in an appropriate metric, with respect to which X_S is compact and the group $\mathbb{R}^d$ acts continuously on X_S by translations, defining a *substitution dynamical system*. The ergodic and spectral properties of such dynamical systems were studied in detail by Solomyak [14].

A substitution tiling space X_S is then associated to an hierarchy. The zero level of this hierarchy is constituted by the initial set of tiles and the level $i > 0$ is constituted by the patches of tiles obtained from those of level $i - 1$ by applying the substitution map S. Recently, Frank and Sadun [4] have introduced a framework to handling with general hierarchical (*fusion*) tiling spaces, where the procedure for obtaining patches of level i from those of level $i-1$ is not necessarily an "inflate-subdivide" procedure and can depend on i. However, many of the ergodic and spectral properties available for substitution dynamical systems are hard to achieve in such generality.

In the present paper we deal with *multi-substitution tiling spaces*, also referred to in the literature as *mixed substitution* tiling spaces [5] or *S-adic systems* [1, 2]. They form a particular class of hierarchical tiling spaces which includes the substitution tiling spaces. A multi-substitution tiling space is determined by a finite number (up to translation) of tiles, a finite set $\mathcal{S} = \{S_1 \ldots, S_k\}$ of substitutions maps acting on these tiles and a sequence

Date: July 17, 2012.

2010 *Mathematics Subject Classification.* 37A15, 37A25, 52C22.

Key words and phrases. multi-substitutions, tiling spaces, dynamical systems, invariant measures, statistical stability.

The authors were partially supported by the Portuguese Government through FCT, under the project PEst-OE/MAT/UI0212/2011 (CMUBI).

$\bar{a} = (a_1, a_2, \ldots)$ in $\Sigma := \{1, \ldots, k\}^{\mathbb{N}}$. In the corresponding hierarchy, the patches of the level i are obtained from those of level $i-1$ by applying the substitution map S_{a_i}. The continuous action of $\mathbb{R}^d$ by translations on a multi-substitution tiling space $X_{\bar{a}}(\mathcal{S})$ defines a uniquely ergodic dynamical system. The unique measure $\mu_{\bar{a},\mathcal{S}}$ is closely related with the patch frequencies in tilings of $X_{\bar{a}}(\mathcal{S})$. In this paper we prove that, in the usual topology of Σ, the ergodic limits of patch frequencies vary continuously with $\bar{a}$ (theorem 20). Moreover, we prove that the convergence of patch frequencies to their ergodic limits is locally uniform in some open subset of Σ (theorem 18).

2. Tilings and Substitutions

We start by recalling some standard definitions and results concerning substitution tiling spaces. For details, motivation and examples see [3, 8, 12, 14]. We introduce also the concept of *strongly recognizable* substitution. As we will see later, such substitutions provide isomorphisms between ergodic dynamical systems associated to certain multi-substitution tiling spaces.

Consider $\mathbb{R}^d$ with its usual Euclidean norm $\|\cdot\|$ and write $B_r = \{\vec{v} \in \mathbb{R}^d : \|\vec{v}\| \leq r\}$. A set $D \subset \mathbb{R}^d$ is called a *tile* if it is compact, connected and equal to the closure of its interior. A *patch* is a collection $x = \{D_i\}_{i\in I}$ of tiles such that $D_i^\circ \cap D_j^\circ = \emptyset$, for all $i, j \in I$ with $i \neq j$. The *support* of x is defined by $\mathrm{supp}(x) := \bigcup_{i\in I} D_i$. If $\mathrm{supp}(x) = \mathbb{R}^d$, we say that x is a *tiling* of $\mathbb{R}^d$. When a patch has a single tile D, we identify this patch with the corresponding tile. Given a patch $x = \{D_i\}_{i\in I}$ and $\vec{t} \in \mathbb{R}^d$, $\vec{t} + x := \{\vec{t} + D_i\}_{i\in I}$ is another patch. In particular, if x is a tiling of $\mathbb{R}^d$, $\vec{t} + x$ is another tiling of $\mathbb{R}^d$. Hence we have an action of $\mathbb{R}^d$ on the space of all tilings of $\mathbb{R}^d$ by translations, which we denote by T. Two patches x and x' are said to be *equivalent* if $x' = \vec{t} + x$ for some $\vec{t} \in \mathbb{R}^d$. We denote by $[x]$ the equivalence class of x.

Let X be a space of tilings of $\mathbb{R}^d$ invariant by T and $\mathcal{P}^N(X)$ be the set of all patches $x' = \{D_i\}_{i\in I}$ such that $|I| = N$ and $x' \subset x$ for some $x \in X$. We denote by $\mathcal{T}^N(X)$ the set of equivalence classes with representatives in $\mathcal{P}^N(X)$. These representatives are called *N-protopatches* of X. The 1-protopatches are more usually called *prototiles*. The tiling space X has *finite local complexity* if $\mathcal{T}^2(X)$ is finite. Equivalently, $\mathcal{T}^N(X)$ is finite for each N.

If $K \subset \mathbb{R}^d$ is compact and $x \in X$, we denote by $x[[K]]$ the set of all patches $x' \subset x$ with bounded support satisfying $K \subseteq \mathrm{supp}(x')$. For $x, y \in X$, we set

$$d_T(x,y) = \inf\Big\{\{\sqrt{2}/2\} \cup \{0 < r < \sqrt{2}/2 : \text{ exist } x' \in x[[B_{1/r}]],\ y' \in y[[B_{1/r}]],\\ \text{and } \vec{t} \in \mathbb{R}^d \text{ with } \|\vec{t}\| \leq r \text{ and } \vec{t} + x' = y'\}\Big\}.$$

Theorem 1. [12, 14] (X, d_T) is a complete metric space. Moreover, if X has finite local complexity, then (X, d_T) is compact and the action T is continuous.

From now on we assume that X is equipped with the metric d_T and that X has finite local complexity.

Remark 1. The above equivalence relation between patches and the corresponding definition of distance could be defined with respect to rather general "actions" of groups on patches (see [10]). However, the metric d_T is adequate to the purposes of this paper since we shall only be concerned with the dynamics associated to the action T.

A *(self-similar) substitution* is a map $S : \mathcal{P}^1(X) \to \mathcal{P}(X) := \bigcup_N \mathcal{P}^N(X)$ such that:

(S_1) there is $\lambda > 1$ (the *dilatation factor* of S) such that $\text{supp}(S(P)) = \lambda \text{supp}(P)$ for all $P \in \mathcal{P}^1(X)$;

(S_2) if $P = \vec{t} + Q$ then $S(P) = \lambda\vec{t} + S(Q)$.

Take a finite number of (non-equivalent) prototiles $\{D_1, \ldots, D_l\}$ of X such that

$$\mathcal{T}^1(X) = \{[D_1], \ldots, [D_l]\}.$$

The *structure matrix* A_S associated to the substitution S is the $l \times l$ matrix with entries A_{ij} equal to the number of tiles equivalent to D_i that appear in $S(D_j)$. If $A_S^m > 0$ for some $m > 0$, S is said to be *primitive*. In the particular case $m = 1$, S is *strongly primitive*.

Given a patch $x = \{D_i\}_{i\in I}$ with $D_i \in \mathcal{P}^1(X)$, we define the patch $S(x) := \bigcup_{i\in I} S(D_i)$. Assume that the substitution S can be extended to maps $S : \mathcal{P}(X) \to \mathcal{P}(X)$ and $S : X \to X$. In this case, take a tile $D \in \mathcal{P}^1(X)$ and define inductively the following sequence of patches in $\mathcal{P}(X)$: $x_1 = D$, and $x_k = S(x_{k-1})$ for $k > 1$. Consider the closed (hence compact) tiling space $X_S \subseteq X$, commonly known as *substitution tiling space* associated to S, defined by: a tiling $x \in X$ belongs to X_S if, and only if, for any finite patch $x' \subset x$ there exist $k > 0$ and a vector $\vec{t} \in \mathbb{R}^d$ such that $\vec{t} + x' \subseteq x_k$. We have:

Proposition 2. [12, 14] Suppose that S is primitive. Then $X_S \neq \emptyset$, $S(X_S) \subseteq X_S$ and X_S is independent of the initial tile $D \in \mathcal{P}^1(X)$.

The Perron-Frobenius (PF) theorem for non-negative matrices is a crucial tool for the study of substitution tiling spaces:

Theorem 3. [13] Let $A \geq 0$ be a real square matrix with $A^m > 0$ for some $m > 0$. Then there is a simple positive eigenvalue $\omega > 0$ of A with $\omega > |\omega'|$ for all other eigenvalues ω'. Moreover, there exist eigenvectors $\vec{p}$ and $\vec{q}$ corresponding to ω for A and A^T, respectively, such that $\vec{p} \cdot \vec{q} = 1$ and $\vec{p}, \vec{q} > 0$. In this case, for any $\vec{v} \in \mathbb{R}^d$

$$\lim_n \frac{A^n\vec{v}}{\omega^n} = (\vec{q} \cdot \vec{v})\vec{p}.$$

The eigenvalue $\omega > 0$ is called the *PF-eigenvalue* of A. The eigenvectors $\vec{p} > 0$ and $\vec{q} > 0$ are called the *right PF-eigenvector* and *left PF-eigenvector* of A, respectively.

In general, if S is a substitution acting on a set of prototiles $\{D_1, \ldots, D_l\}$ with Euclidian volumes $V_1, \ldots, V_l$, the vector $\vec{q} = (V_1, \ldots, V_l)$ is a left eigenvector of A_S associated to the eigenvalue λ^d. For primitive substitutions, $\vec{q}$ and $\omega = \lambda^d$ are precisely the left PF-eigenvector and the corresponding PF-eigenvalue of A_S, respectively.

A substitution S is said to be *recognizable* if $S : X \to S(X)$ is injective. In this case, we say that S is *strongly recognizable* if for any $x \in X$ and any tile $D \in \mathcal{P}^1(X)$ the following holds: if $S(D)$ is a patch of $S(x)$, then $D \in x$.

Example 1. The Ammann A3 substitution (figure 1; see [8] for a detailed description of this substitution) is recognizable but not strongly recognizable. More generally, any sub-




FIGURE 1. Ammann A3 substitution.

stitution for which one patch $S(D_i)$ contains another patch $S(D_j)$ can not be strongly recognizable. The pentiamond substitution (figure 2; see the Tilings Encyclopedia at http://tilings.math.uni-bielefeld.de) is strongly recognizable.



FIGURE 2. Pentiamond substitution.

3. Multi-Substitution Tiling Spaces

In this section we establish the definition and basic properties of multi-substitution tiling spaces. These spaces are also referred to as *mixed substitution* tiling spaces [5] or *S-adic systems* [1, 2]. In [4], the authors developed a framework for studying the ergodic theory and topology of hierarchical tilings in great generality. The classical substitution tiling spaces and the multi-substitution tiling spaces fit in this general framework. In fact, they are particular cases of *fusion* tiling spaces. Certain properties, like minimality or unique ergodicity, can be derived within the framework of fusion tiling spaces. However, naturally, some other properties become hard to achieve in such generality.

Let $\mathcal{S} = \{S_i\}_{i\in J}$ be a finite collection of substitutions $S_i : \mathcal{P}^1(X) \to \mathcal{P}(X)$. Assume that the substitution S_i can be extended to maps $S_i : \mathcal{P}(X) \to \mathcal{P}(X)$ and $S_i : X \to X$, for each $i \in J$. Denote by $\lambda_i > 1$ and A_i the dilatation factor and the structure matrix, respectively, associated to S_i. Provide the space of sequences

$$\Sigma := \{\bar{a} = (a_1, a_2, \ldots) : a_i \in J\}$$

with the usual structure (Σ, d_Σ) of metric space: given $\bar{a} = (a_1, a_2, \ldots)$ and $\bar{b} = (b_1, b_2, \ldots)$ in Σ, we set $d_\Sigma(\bar{a}, \bar{b}) = 1/L$ if L is the smallest integer such that $a_L \neq b_L$. We also introduce the standard shift map $\sigma : \Sigma \to \Sigma$, given by $\sigma(a_1, a_2, \ldots) = (a_2, a_3, \ldots)$, which is continuous with respect to d_Σ. Given $\bar{a} = (a_1, a_2, \ldots) \in \Sigma$, we denote by $[\bar{a}]_n$ the periodic sequence

$(a_1, ..., a_n, a_1, ..., a_n, \ldots) \in \Sigma$. Clearly, for each $k > 0$, $S_{\bar{a}}^k := S_{a_1} \circ S_{a_2} \circ \ldots \circ S_{a_k}$ is itself a substitution with structure matrix given by $A_{\bar{a}}^k := A_{a_1} A_{a_2} \ldots A_{a_k}$ and dilation factor $\lambda_{\bar{a}}^n := \lambda_{a_1} \lambda_{a_2} \ldots \lambda_{a_n}$.

The sequence of substitutions (S_{a_n}) is called *primitive* if for each n there exists a least $N_n^{\bar{a}}$ such that $A_{a_n} A_{a_{n+1}} \ldots A_{a_{n+N_n^{\bar{a}}}} > 0$. Observe that, in this case, each matrix A_i does not have any column of all zeroes. Hence, $A_{a_n} A_{a_{n+1}} \ldots A_{a_{n+N_n^{\bar{a}}+j}} > 0$ for all $j \geq 0$. If, for each n, the substitution S_{a_n} is strongly primitive, that is, $A_{a_n} > 0$, the sequence (S_{a_n}) is called *strongly primitive*. The set $\mathcal{S}$ of substitutions is *primitive* if, for any $\bar{a} \in \Sigma$, (S_{a_n}) is primitive. Moreover, we say that a primitive set of substitutions $\mathcal{S}$ is *bounded primitive* if the set $\{N_n^{\bar{a}} : n \in \mathbb{N}, \bar{a} \in \Sigma\}$ is bounded.

Lemma 4. If $\mathcal{S}$ is primitive then it is bounded primitive.

Proof. The number of possible configurations of zero entries in finite products of structure matrices associated to substitutions in $\mathcal{S}$ is finite, say $L(\mathcal{S})$. Now, assume that $\mathcal{S}$ is not bounded primitive. Then, for some $q > L(\mathcal{S})$, there is $\bar{a} \in \Sigma$ such that $A_{\bar{a}}^q$ has some zero entry. This means that we can find $p' < p \leq q$ such that $A_{\bar{a}}^{p'}$ has the same zero configuration of entries as $A_{\bar{a}}^p$. Then the sequence of substitutions (S_{b_n}), with $\bar{b} = (a_1, \ldots, a_{p'}, a_{p'+1}, \ldots, a_p, a_{p'+1}, \ldots, a_p, a_{p'+1}, \ldots)$ is non-primitive. □

Before introduce the multi-substitution tiling spaces, let us prove the following useful lemma.

Lemma 5. Take $D \in \mathcal{P}^1(X)$ and assume that the sequence of substitutions (S_{a_n}) is primitive. Given $n > 0$, there are $R > 0$ and $N_0 > n$ such that, for any $N > N_0$ and any ball B of radius R contained in the support of $S_{\bar{a}}^N(D)$, the following holds: $\vec{t} + S_{\bar{a}}^n(D) \subset S_{\bar{a}}^N(D)$ and $\text{supp}(\vec{t} + S_{\bar{a}}^n(D)) \subset B$, for some $\vec{t} \in \mathbb{R}^d$.

Proof. Take a finite number of prototiles $\{D_1, \ldots, D_l\}$ of X such that

$$\mathcal{T}^1(X) = \{[D_1], \ldots, [D_l]\}.$$

By primitivity, we can take N_0 such that a translated copy of $S_{\bar{a}}^n(D)$ can be found in each $S_{\bar{a}}^{N_0}(D_i)$ for all $i \in \{1, \ldots, l\}$. Now, consider the tiles $\tilde{D}_i = \text{supp}(S_{\bar{a}}^{N_0}(D_i))$. For sufficiently large R, if B is a ball with radius R and x' is a patch formed with translated copies of the tiles $\tilde{D}_i$, with $B \subset \text{supp}(x')$, then, for some $i \in \{1, \ldots, l\}$ and $\vec{t}_i \in \mathbb{R}^d$, we have $\vec{t}_i + \tilde{D}_i \subset B$ and $\vec{t}_i + \tilde{D}_i \in x'$. Take $N > N_0$ such that the support of $S_{\bar{a}}^N(D)$ contains some ball B of radius R. Since $S_{\bar{a}}^N(D)$ is the disjoint union of translated copies of patches of the form $S_{\bar{a}}^{N_0}(D_i)$, the support of one of this copies must be contained in B, and we are done. □

Remark 2. It is clear from the proof that, given $n > 0$ and tile D, if $\mathcal{S}$ is primitive (hence bounded primitive) we can take $R > 0$ and $N_0 > n$ so that the statement of lemma 5 holds for any $\bar{a} \in \Sigma$.

Take $D \in \mathcal{P}^1(X)$, $\bar{a} \in \Sigma$ and the corresponding sequence of patches in $\mathcal{P}(X)$:

$$x_{\bar{a}}^1 = D, \quad \text{and} \quad x_{\bar{a}}^k = S_{\bar{a}}^k(D), \text{ for } k > 1.$$

We define the *multi-substitution tiling space* $X_{\bar{a}} := X_{\bar{a}}(\mathcal{S}) \subseteq X$ as follows: a tiling $x \in X$ belongs to $X_{\bar{a}}$ if, and only if, for any finite patch $x' \subset x$ there exist $k > 0$ and a vector $\vec{t} \in \mathbb{R}^d$ such that $\vec{t} + x' \subseteq x_{\bar{a}}^k$.

Proposition 6. If the sequence of substitutions (S_{a_n}) is primitive, then:

a) $X_{\bar{a}} \neq \emptyset$;
b) $X_{\bar{a}}$ is independent of the initial tile $D \in \mathcal{P}^1(X)$;
c) $X_{[\bar{a}]_n}$ coincides with the substitution tiling space $X_{S^n_{\bar{a}}}$;
d) $X_{\bar{a}}$ is closed.

Proof. Take $D \in \mathcal{P}^1(X)$. By primitivity, the definition of $X_{\bar{a}}$ is independent of the initial tile and, taking account lemma 5, for each $n > 0$ there are $\vec{t}_n \in \mathbb{R}^d$, $k_n > 0$ and $r_n > 0$, with $\lim_n r_n = \infty$, such that the sequence (x'_n) of patches $x'_n = \vec{t}_n + S^{k_n}_{\bar{a}}(D)$ satisfies: $x'_{n-1} \subset x'_n$ and $\mathrm{supp}(x'_{n-1}) \subset B_{r_n} \subset \mathrm{supp}(x'_n)$. Set

$$x^{\infty}_{\bar{a}} = \bigcup_{n\geq 1} x'_n \tag{1}$$

and observe that $x^{\infty}_{\bar{a}}$ is a tiling of $\mathbb{R}^d$ in $X_{\bar{a}}$. Hence $X_{\bar{a}} \neq \emptyset$.

It is clear that $X_{S^n_{\bar{a}}} \subset X_{[\bar{a}]_n}$. Now, take $x \in X_{[\bar{a}]_n}$ and a finite patch $x' \subset x$. By definition, a translated copy of x' appears in $S^k_{[\bar{a}]_n}(D)$ for some $k > 0$. Take $N > k$ such that $A_{a_{k+1}} \dots A_{a_N} > 0$ and m such that $N \leq mn$. Then we also have $A_{a_{k+1}} \dots A_{a_{nm}} > 0$. In particular, a translated copy of D appears in $S_{a_{k+1}} \circ \dots \circ S_{a_{nm}}(D)$. Consequently, a translated copy of x' appears in $S^{nm}_{[\bar{a}]_n}(D) = (S^n_{[\bar{a}]_n})^m(D)$. This means that $x \in X_{S^n_{\bar{a}}}$.

To prove that $X_{\bar{a}}$ is closed, take a sequence of tilings (x_n) in $X_{\bar{a}}$ converging to some $x \in X$. Take a patch x'' in x and $r > 0$ such that $\mathrm{supp}(x'') \subset B_{1/r}$. We know that there exists n_0 such that $d_T(x_n, x) < r$ for all $n > n_0$. This means that, for each $n > n_0$, there are patches $x'_n \in x_n[[B_{1/r}]]$ and $x' \in x[[B_{1/r}]]$, and a vector $\vec{t}_n$ with $\|\vec{t}_n\| < r$, such that $x' = \vec{t}_n + x'_n$. Since $x_n \in X_{\bar{a}}$, there exists a translation of x'_n, and consequently of $x'' \subset x'$, that is contained in some $S^{k_n}_{\bar{a}}(D)$. Hence $x \in X_{\bar{a}}$. □

Henceforth we assume that the sequence of substitutions (S_{a_n}) is primitive. As for substitution tiling spaces, recognizability is closely related with non-periodicity:

Proposition 7. Let $\mathcal{S} = \{S_1, \dots, S_k\}$ be a set of recognizable substitutions, $\bar{a} \in \Sigma$, and $X_{\bar{a}}$ the corresponding multi-substitution tiling space. Then any tiling $x \in X_{\bar{a}}$ is aperiodic.

Proof. The argument is standard. Take $x \in X_{\bar{a}}$ and suppose that $\vec{t} + x = x$ for some $\vec{t} \neq 0$. Since our substitutions are recognizable, for each $n \geq 1$ there exists a unique $x_n \in X_{\sigma^n(\bar{a})}$ such that $S^n_{\bar{a}}(x_n) = x$. We have

$$S^n_{\bar{a}}(x_n) = \vec{t} + S^n_{\bar{a}}(x_n) = S^n_{\bar{a}}\Big(\frac{\vec{t}}{\lambda^n_{\bar{a}}} + x_n\Big),$$

which means, by recognizability that $x_n = \vec{t}/\lambda^n_{\bar{a}} + x_n$. Now, for n sufficiently large, it is clear that $\big\{\mathrm{supp}(D) + \vec{t}/\lambda^n_{\bar{a}}\big\} \cap \mathrm{supp}(D) \neq \emptyset$ for any prototile D, which is a contradiction. □

It is well known that the set of periodic points of σ is dense in Σ. Together with proposition 6, this result suggests that any multi-substitution tiling can be approximated arbitrarily closely by substitution tilings. In fact we have:

Proposition 8. For each $x \in X_{\bar{a}}$, there exists a sequence of tilings (x_n), with $x_n \in X_{S_{\bar{a}}^{j_n}}$, for some $j_n \geq 1$, such that $\lim_n x_n = x$.

Proof. For each n take a patch $x'_n \in x[[B_n]]$. By definition of multi-substitution tiling space, we have $x'_n \subset \vec{t}_n + S_{\bar{a}}^{j_n}(D)$, for some $j_n \geq 1$ and $\vec{t}_n \in \mathbb{R}^d$. Adapting the procedure we have used in the proof of proposition 6 to construct a tiling in $X_{\overline{a}}$, it is possible to construct a tiling $x_n \in X_{S_{\bar{a}}^{j_n}} = X_{[\bar{a}]_{j_n}}$ containing $\vec{t}_n + S_{\bar{a}}^{j_n}(D)$. Clearly we have $\lim_n x_n = x$. □

4. Minimality and Repetitivity

As Sadun and Frank [4] have shown, fusion tiling spaces are minimal and its elements are repetitive. For completeness, we shall next give a proof of this result in the particular case of multi-substitution tiling spaces.

A *dynamical system* will be a pair (Y, G) where Y is a compact metric space and G is a continuous action of a group. (Y, G) is *minimal* if Y is the orbit closure $\overline{\mathcal{O}(y)}$ of any of its elements y. A point $y \in Y$ is *almost periodic* if

$$G(y, U) = \{g \in G : \ g(y) \in U\}$$

is *relatively dense* (that is, there exists a compact set $K \subseteq G$ such that $g \cdot K$ intersects $G(y, U)$ for all $g \in G$) for every open set $U \subseteq Y$ with $G(y, U) \neq \emptyset$. Minimality and almost periodicity are related by Gottschalk's theorem:

Theorem 9. [7] Let (Y, G) be a dynamical system. If $y \in Y$ is an almost periodic point, then $(\overline{\mathcal{O}(y)}, G)$ is minimal. Moreover, if (Y, G) is minimal, then any point in Y is almost periodic.

It is well known [12] that, for any primitive substitution S, (X_S, T) is minimal. More generally, for multi-substitutions tiling spaces we have:

Theorem 10. The dynamical system $(X_{\bar{a}}, T)$ is minimal.

Proof. Let x and y be two tilings in $X_{\bar{a}}$ and $\epsilon > 0$. Fix $y' \in y[[B_{1/\epsilon}]]$. By definition of $X_{\bar{a}}$, there is n such that a translated copy of y' can be found in $S_{\bar{a}}^n(D)$. Taking account lemma 5, there are $R > 0$ and $N' > 0$ such that, for any ball B of radius R with $B \subset \text{supp}(S_{\bar{a}}^{N'}(D))$, there is $\vec{t}$ for which $\vec{t} + S_{\bar{a}}^n(D) \subset S_{\bar{a}}^{N'}(D)$ and $\text{supp}(\vec{t} + S_{\bar{a}}^n(D)) \subset B$. On the other hand, again by definition of $X_{\bar{a}}$, given a patch $x' \in x[[B_R]]$, there is some N'' such that $S_{\bar{a}}^{N''}(D)$ contains a translated copy of x'. Hence, due to primitivity, we can take some $N \geq \max\{N', N''\}$ such that

$$y' \subseteq \vec{t}_1 + S_{\bar{a}}^n(D) \subseteq \vec{t}_2 + x' \subseteq \vec{t}_3 + S_{\bar{a}}^N(D).$$

In particular, $d_T(\vec{t}_2 + x, y) < \epsilon$. □

A tiling x of $\mathbb{R}^d$ is said to be *repetitive* if for every patch x' of x with bounded support there is some $r(x') > 0$ such that, for every ball B of $\mathbb{R}^d$ with radius $r(x')$, there exists $\vec{t} \in \mathbb{R}^d$ such that $\text{supp}(\vec{t} + x') \subseteq B$ and $\vec{t} + x' \subset x$. It is also common to refer to repetitive tilings as tilings satisfying the *local isomorphism* property [11]. As explained in [12], for tiling dynamical systems (X, T), repetitivity is equivalent to almost periodicity. From Gottschalk's theorem it follows that:

Theorem 11. Any $x \in X_{\bar{a}}$ is repetitive.

Of course, this can also be seen as an easy consequence of lemma 5. Given a tiling $x \in X_{\bar{a}}$ and a finite patch $x' \subset x$, we have $x' \subset S^n_{\bar{a}}(D)$ for some n and D. The radius $r(x')$ can be taken as the radius R of lemma 5, which does not depend on the tiling x of $X_{\bar{a}}$ we take. On the other hand, $x' \in \mathcal{P}(X_{\bar{b}})$ for any $\bar{b} \in \Sigma$ with $d_\Sigma(\bar{a}, \bar{b}) < 1/n$. Hence, taking account remark 2, we have:

Proposition 12. Assume that $\mathcal{S}$ is primitive and take $x' \subset S^n_{\bar{a}}(D)$. Then, there exists $R > 0$ such that, for any $\bar{b} \in \Sigma$ with $d_\Sigma(\bar{a}, \bar{b}) < 1/n$, any $x \in X_{\bar{b}}$ and any ball B of radius R, a translated copy $\vec{t} + x'$ of x' can be founded in x with $\mathrm{supp}(\vec{t} + x') \subseteq B$.

5. Statistical stability

5.1. **Unique ergodicity.** The unique ergodicity of the system $(X_{\bar{a}}, T)$ was established in [4] in the general framework of fusion tiling spaces. Next we remake the proof of the unique ergodicity for multi-substitution tiling spaces, based on a result of Solomyak [14], and prove that the convergence of patch frequencies to their ergodic limits is locally uniform in some open subset of Σ (theorem 18).

Given a patch $x' \in \mathcal{P}(X_{\bar{a}})$ and a measurable subset U of $\mathbb{R}^d$, define the *cylinder set* $X^{\bar{a}}_{x',U}$ as

$$X^{\bar{a}}_{x',U} = \{x \in X_{\bar{a}} : \ x' + \vec{t} \subset x \text{ for some } \vec{t} \in U\}.$$

These cylinders form a semi-algebra and a topology base for $X_{\bar{a}}$.

For any set $H \subset \mathbb{R}^d$ and $r \geq 0$ we define

$$H^{+r} = \{\vec{t} \in \mathbb{R}^d : \mathrm{dist}(\vec{t}, H) \leq r\}, \quad H^{-r} = \{\vec{t} \in H : \mathrm{dist}(\vec{t}, \partial H) \geq r\},$$

where ∂H denotes the boundary of H. A sequence (H_n) of subsets of $\mathbb{R}^d$ is a *Van Hove* sequence if for any $r \geq 0$

$$\lim_n \frac{\mathrm{vol}((\partial H_n)^{+r})}{\mathrm{vol}(H_n)} = 0.$$

Clearly, the sequence $(\mathrm{supp}(S^n_{\bar{a}}(D)))$ is a Van Hove sequence for each $D \in \mathcal{P}^1(X)$.

Consider the tiling $x^\infty_{\bar{a}}$ given by (1). For any patch $x' \in \mathcal{P}(X_{\bar{a}})$, denote by $L^{\bar{a}}_{x'}(H)$ (respectively, $N^{\bar{a}}_{x'}(H)$) the number of distinct translated copies of x' in $x^\infty_{\bar{a}}$ whose support is completely contained in H (respectively, intersects the border of H). If $y' \in \mathcal{P}(X_{\bar{a}})$ is another patch, we denote by $L_{x'}(y')$ the number of distinct translated copies of x' in y' and by $\mathrm{vol}(y')$ the Euclidean volume of the support of y'.

Theorem 13. [14] The dynamical system $(X_{\bar{a}}, T)$ is uniquely ergodic if for any patch $x' \in \mathcal{P}(X_{\bar{a}})$ there is a number $\mathrm{freq}_{\bar{a}}(x') > 0$ such that, for any Van Hove sequence (H_n),

$$\mathrm{freq}_{\bar{a}}(x') = \lim_n \frac{L^{\bar{a}}_{x'}(H_n)}{\mathrm{vol}(H_n)}.$$

In this case, the unique ergodic measure $\mu_{\bar{a}}$ on $X_{\bar{a}}$ satisfies

$$\mu_{\bar{a}}(X^{\bar{a}}_{x',U}) = \mathrm{freq}_{\bar{a}}(x')\mathrm{vol}(U)$$

for all Borel subsets U with $\mathrm{diam}(U) < \eta$, where $\eta > 0$ is such that any prototile contains a ball of radius η.

For a primitive substitution tiling space X_S, such *frequencies* exist [6, 14]. For example, if $\vec{p} = (p_1, \dots, p_l)$ and $\vec{q} = (V_1, \dots, V_l)$ are right and left PF-eigenvectors, respectively, of A_S satisfying $\vec{p} \cdot \vec{q} = 1$, then $\mathrm{freq}_S(D_i) = p_i$. Consequently, (X_S, T) is uniquely ergodic. We want to extend this result for multi-substitutions tiling spaces.

So, start with a primitive set $\mathcal{S} = \{S_1, \dots, S_k\}$ of substitutions acting on the set of prototiles $\{D_1, \dots, D_l\}$. Being primitive, there exists some $N(\mathcal{S}) > 0$ such that $A > 0$ for any $A \in \mathcal{A}_{\mathcal{S}}^{N(\mathcal{S})}$, the set of all $N(\mathcal{S})$-products of matrices from $\{\frac{A_1}{\omega_1}, \dots, \frac{A_k}{\omega_k}\}$, where ω_i is the PF-eigenvalue of A_i.

For each $i, j \in \{1, \dots, l\}$, set

$$c_{\bar{a}}(D_i) := \lim_n \frac{L_{D_i}(S_{\bar{a}}^n(D_j))}{\mathrm{vol}(S_{\bar{a}}^n(D_j)))} = \lim_n \frac{(A_{\bar{a}}^n)_{ij}}{\omega_{\bar{a}}^n \mathrm{vol}(D_j)}. \tag{2}$$

Lemma 14. The limit (2) exists, does not depend on j and is uniform with respect to $\bar{a} \in \Sigma$. Moreover, there are $C > 0$ and $0 < \theta < 1$ such that, for any $\delta > 0$ and $\bar{a}, \bar{b} \in \Sigma$ with $d_\Sigma(\bar{a}, \bar{b}) < \delta$, we have

$$|c_{\bar{a}}(D_i) - c_{\bar{b}}(D_i)| < C\theta^{1/\delta}, \tag{3}$$

for all $i \in \{1, \dots, l\}$.

Proof. Let $\bar{a} \in \Sigma$. For each $n \geq 1$, consider the matrix $E_{\bar{a}}^n = [(E_{\bar{a}}^n)_{ij}]$ defined by

$$(E_{\bar{a}}^n)_{ij} = \frac{(A_{\bar{a}}^n)_{ij}}{\omega_{\bar{a}}^n \mathrm{vol}(D_j)}.$$

Let $\Delta_{\bar{a}}^n \subset \mathbb{R}^d$ be the convex hull of the columns of $E_{\bar{a}}^n$. It is easy to check that each column of $E_{\bar{a}}^{n+N}$ sits in $\Delta_{\bar{a}}^n$. Indeed,

$$(E_{\bar{a}}^{n+N})_{ij} = \sum_{k=1}^{l} \frac{(A_{\bar{a}}^n)_{ik}}{\omega_{\bar{a}}^n \mathrm{vol}(D_k)} \frac{(A_{\sigma^n(\bar{a})}^N)_{kj} \mathrm{vol}(D_k)}{\omega_{\sigma^n(\bar{a})}^N \mathrm{vol}(D_j)}.$$

Hence, the j-column $v_{\bar{a},j}^{n+N}$ of $E_{\bar{a}}^{n+N}$ is given by

$$v_{\bar{a},j}^{n+N} = \sum_{k=1}^{l} v_{\bar{a},k}^n \frac{(A_{\sigma^n(\bar{a})}^N)_{kj} \mathrm{vol}(D_k)}{\omega_{\sigma^n(\bar{a})}^N \mathrm{vol}(D_j)}. \tag{4}$$

Since

$$\sum_{k=1}^{l} \frac{(A_{\sigma^n(\bar{a})}^N)_{kj} \mathrm{vol}(D_k)}{\omega_{\sigma^n(\bar{a})}^N \mathrm{vol}(D_j)} = 1,$$

we have $v_{\bar{a},j}^{n+N} \in \Delta_{\bar{a}}^n$, that is $\Delta_{\bar{a}}^{n+N} \subseteq \Delta_{\bar{a}}^n$ for all $N > 0$. Set $\Delta_{\bar{a}} = \bigcap_{n=1}^{\infty} \Delta_{\bar{a}}^n$ and take θ' (not depending on $\bar{a}$) satisfying $\theta' l < 1$ and

$$0 < \theta' < \min_{A \in \mathcal{A}_{\mathcal{S}}^{N(\mathcal{S})}} \min_{k,j} \Big\{ \frac{A_{kj} \mathrm{vol}(D_k)}{\mathrm{vol}(D_j)} \Big\}.$$

Taking account (4), we have

$$v_{\bar{a},j}^{n+N(\mathcal{S})} = \sum_{k=1}^{l} (1 - \theta' l) v_{\bar{a},k}^n \Big(\frac{(A_{\sigma^n(\bar{a})}^{N(\mathcal{S})})_{kj} \mathrm{vol}(D_k)}{(1 - \theta' l) \omega_{\sigma^n(\bar{a})}^{N(\mathcal{S})} \mathrm{vol}(D_j)} - \frac{\theta'}{1 - \theta' l} \Big) + \sum_{k=1}^{l} v_{\bar{a},k}^n \theta'.$$

Observe that

$$\sum_{k=1}^{l}\Big(\frac{(A^{N(\mathcal{S})}_{\sigma^n(\bar{a})})_{kj}\mathrm{vol}(D_k)}{(1-\theta' l)\omega^{N(\mathcal{S})}_{\sigma^n(\bar{a})}\mathrm{vol}(D_j)}-\frac{\theta'}{1-\theta' l}\Big)=1.$$

Hence $\Delta^{n+N(\mathcal{S})}_{\bar{a}}$ is contained, up to translation, in $(1-\theta' l)\Delta^n_{\bar{a}}$. Then

$$\mathrm{diam}(\Delta^{1+nN(\mathcal{S})}_{\bar{a}})\leq\rho(1-\theta' l)^n \tag{5}$$

for all n, where ρ is the maximum of the l possible values of $\mathrm{diam}(\Delta^1_{\bar{a}})$. Consequently $\Delta_{\bar{a}}$ has diameter zero, hence it consists of a single point, which means that the limit (2) exists and does not depend on j. Moreover, this limit is uniform with respect to $\bar{a}\in\Sigma$ since the majoration (5) does not depend on $\bar{a}$.

Finally, note that if $d_\Sigma(\bar{a},\bar{b})=1/k<\delta$, that is $a_j=b_j$ and $\Delta_j:=\Delta^j_{\bar{a}}=\Delta^j_{\bar{b}}$ for all $j\in\{1,\dots,k-1\}$, then

$$|c_{\bar{a}}(D_i)-c_{\bar{b}}(D_i)|\leq\mathrm{diam}(\Delta_{k-1})\leq C\theta^{1/\delta}$$

for $\theta=(1-\theta' l)^{1/N(\mathcal{S})}$ and some constant $C>0$. □

Take a finite patch $x'\in\mathcal{P}(X)$. For $\bar{a}\in\Sigma$, set

$$c_{\bar{a}}(x'):=\lim_n\frac{L_{x'}(S^n_{\bar{a}}(D_j))}{\mathrm{vol}(S^n_{\bar{a}}(D_j)))}. \tag{6}$$

Lemma 15. The limit (6) exists and does not depend on j. Moreover, if $x'\in\mathcal{P}(X_{\bar{a}})$, then the limit is uniform in a small neighborhood of $\bar{a}$.

Proof. Take a finite patch $x'\in\mathcal{P}(X)$ and a sequence $\bar{a}\in\Sigma$. Given a subset H of $\mathbb{R}^d$, observe that

$$\frac{L^{\bar{a}}_{x'}(H)}{\mathrm{vol}(H)}\leq\frac{1}{\mathrm{vol}(x')}. \tag{7}$$

Given $n>m$, write

$$S^{n-m}_{\sigma^m(\bar{a})}(D_j)=\bigcup_{i,k}D_{ik}\,,$$

where $i\in\{1,\dots,l\}$, $k\in\{1,\dots,\big(A^{n-m}_{\sigma^m(\bar{a})}\big)_{ij}\}$ and each D_{ik} is a translated copy of the prototile D_i. Denote by $N^{\bar{a}}_{x'}(n,m,i,j)$ the number of translated copies of x' contained in $S^n_{\bar{a}}(D_j)$ whose support intersects the boundary of some $\mathrm{supp}(S^m_{\bar{a}}(D_{ik}))$. Since the sequence $(S^m_{\bar{a}}(D_i))$ is Van Hove, for each $\epsilon>0$ there exists $m(\bar{a})$ such that

$$N^{\bar{a}}_{x'}(n,m,i,j)<\epsilon L_{x'}(S^m_{\bar{a}}(D_i))\big(A^{n-m}_{\sigma^m(\bar{a})}\big)_{ij},$$

for all $m>m(\bar{a})$ and $i\in\{1,\dots,l\}$. Hence

$$\sum_{i=1}^{l}\frac{L_{x'}(S^m_{\bar{a}}(D_i))\big(A^{n-m}_{\sigma^m(\bar{a})}\big)_{ij}}{\omega^m_{\bar{a}}\mathrm{vol}(S^{n-m}_{\sigma^m(\bar{a})}(D_j))}\leq\frac{L_{x'}(S^n_{\bar{a}}(D_j))}{\mathrm{vol}(S^n_{\bar{a}}(D_j)))}\leq(1+\epsilon)\sum_{i=1}^{l}\frac{L_{x'}(S^m_{\bar{a}}(D_i))\big(A^{n-m}_{\sigma^m(\bar{a})}\big)_{ij}}{\omega^m_{\bar{a}}\mathrm{vol}(S^{n-m}_{\sigma^m(\bar{a})}(D_j))}. \tag{8}$$

Taking account (7) and lemma 14, we have

$$\limsup_n\frac{L_{x'}(S^n_{\bar{a}}(D_j))}{\mathrm{vol}(S^n_{\bar{a}}(D_j))}-\liminf_n\frac{L_{x'}(S^n_{\bar{a}}(D_j))}{\mathrm{vol}(S^n_{\bar{a}}(D_j))}\leq\frac{\epsilon}{\mathrm{vol}(x')}.$$

Since this holds for an arbitrary ϵ, it follows that the limit $c_{\bar{a}}(x')$ exists. The independence of $c_{\bar{a}}(x')$ with respect to j follows from the observation that the lower and upper bounds in (8) do not depend on j when n goes to infinity.

Suppose now that $x' \in \mathcal{P}(X_{\bar{a}})$. Let us prove that the limit is uniform in a small neighborhood of $\bar{a}$. We have $x' \subset S^{n_0}_{\bar{a}}(D)$ for some n_0 and D. Take the radius $R > 0$ associated to x' given by proposition 12. For $\lambda > 0$, let $L(R, \lambda, i)$ be the maximum number of open disjoint balls of radius R contained in λD_i. Clearly, there exists λ_0 such that

$$\frac{\mathrm{vol}((\partial \lambda D_i)^{+r})}{\mathrm{vol}(x')} < \epsilon L(R, \lambda, i)$$

for all $\lambda > \lambda_0$, with r the diameter of $\mathrm{supp}(x')$. Let $\omega = \min_i\{\omega_i\}$ and take $m_0 > 0$ such that $\omega^{m_0} > \lambda_0$. Then, for all $m > m_0$ and $\bar{b} \in \Sigma$ with $d_\Sigma(\bar{a}, \bar{b}) < 1/n_0$, we have $x' \in \mathcal{P}(X_{\bar{b}})$ and

$$\begin{aligned} N^{\bar{b}}_{x'}(n, m, i, j) &< \frac{\mathrm{vol}\big((\partial \omega^m_{\bar{b}} D_i)^{+r}\big)}{\mathrm{vol}(x')} \big(A^{n-m}_{\sigma^m(\bar{b})}\big)_{ij} \\ &< \epsilon L(R, \lambda, i) \big(A^{n-m}_{\sigma^m(\bar{b})}\big)_{ij} \\ &< \epsilon L_{x'}(S^m_{\bar{b}}(D_i)) \big(A^{n-m}_{\sigma^m(\bar{b})}\big)_{ij}, \end{aligned}$$

and we are done. □

Lemma 16. Assume that the limit (6) exists. Then, for any Van Hove sequence (H_n),

$$c_{\bar{a}}(x') = \lim_n \frac{L^{\bar{a}}_{x'}(H_n)}{\mathrm{vol}(H_n)}.$$

Proof. Although the proof follows closely that of [9] for substitution tilings, we present it here since it is essential to establish theorem 18. For each $m > 0$, there is $\mathcal{D}^{\bar{a}}_m \subseteq \mathcal{P}^1(X_{\bar{a}})$ such that the tiling $x^\infty_{\bar{a}}$ in (1) is the disjoint union of patches $S^m_{\bar{a}}(D)$, with $D \in \mathcal{D}^{\bar{a}}_m$. Given $m > 0$ and $n > 0$, define

$$G^{\bar{a}}_{m,n} := \{D \in \mathcal{D}^{\bar{a}}_m : \mathrm{supp}(S^m_{\bar{a}}(D)) \cap H_n \neq \emptyset\}, \;\; H^{\bar{a}}_{m,n} := \{D \in \mathcal{D}^{\bar{a}}_m : \mathrm{supp}(S^m_{\bar{a}}(D)) \subseteq H_n\}.$$

Hence

$$\sum_{D \in H^{\bar{a}}_{m,n}} L_{x'}(S^m_{\bar{a}}(D)) \leq L^{\bar{a}}_{x'}(H_n) \leq \sum_{D \in G^{\bar{a}}_{m,n}} \Big(L_{x'}(S^m_{\bar{a}}(D)) + N^{\bar{a}}_{x'}(\partial S^m_{\bar{a}}(D)) \Big), \tag{9}$$

where $\partial S^m_{\bar{a}}(D)$ denotes the border of the support of $S^m_{\bar{a}}(D)$. Now, fix $\epsilon > 0$. Taking account that $(S^m_{\bar{a}}(D_j))$ is a Van Hove sequence, we can take m large enough so that, for every $D \in \mathcal{D}^{\bar{a}}_m$, we have

$$\left| \frac{L_{x'}(S^m_{\bar{a}}(D))}{\mathrm{vol}(S^m_{\bar{a}}(D))} - c_{\bar{a}}(x') \right| < \epsilon \quad \text{and} \quad N_{x'}(\partial S^m_{\bar{a}}(D)) < \epsilon L_{x'}(S^m_{\bar{a}}(D)). \tag{10}$$

Together with (9), this gives

$$(c_{\bar{a}}(x') - \epsilon) \sum_{D \in H^{\bar{a}}_{m,n}} \mathrm{vol}(S^m_{\bar{a}}(D)) \leq L^{\bar{a}}_{x'}(H_n) \leq (1 + \epsilon)(c_{\bar{a}}(x') + \epsilon) \sum_{D \in G^{\bar{a}}_{m,n}} \mathrm{vol}(S^m_{\bar{a}}(D)). \tag{11}$$

On the other hand, note that, setting $t_m := \max_j\{\mathrm{diam}(S^m_{\bar{a}}(D_j)\}$, since m is fixed, for large enough n we have

$$\sum_{D\in H^{\bar{a}}_{m,n}} \mathrm{vol}(S^m_{\bar{a}}(D)) \geq \mathrm{vol}(H_n^{-t_m}) \geq (1-\epsilon)\mathrm{vol}(H_n)$$

and

$$\sum_{D\in G^{\bar{a}}_{m,n}} \mathrm{vol}(S^m_{\bar{a}}(D)) \leq \mathrm{vol}(H_n^{+t_m}) \leq (1+\epsilon)\mathrm{vol}(H_n),$$

which, combining with (11), concludes the proof, since $\epsilon > 0$ is arbitrary. □

Combining theorem 13 with lemmas 15 and 16, with $c_{\bar{a}}(x') = \mathrm{freq}_{\bar{a}}(x')$, we conclude that

Theorem 17. If $\mathcal{S}$ is primitive, $(X_{\bar{a}}, T)$ is uniquely ergodic for all $\bar{a} \in \Sigma$.

The following theorem establishes that the patch frequencies converge uniformly to their ergodic limits in an open subset of Σ.

Theorem 18. If $x' \in \mathcal{P}(X_{\bar{a}})$ and H_n is a Van Hove sequence, then the sequences

$$\frac{L^\bullet_{x'}(H_n)}{\mathrm{vol}(H_n)}$$

converge uniformly to $\mathrm{freq}_\bullet(x')$ in a small neighborhood of $\bar{a}$.

Proof. This follows easily from lemma 15 and from the proof of lemma 16, since (10) holds for any sequence in a small neighborhood of $\bar{a}$. □

We denote by $\mu_{\bar{a}} := \mu_{\bar{a},\mathcal{S}}$ the unique ergodic measure of $(X_{\bar{a}}, T)$. Two ergodic dynamical systems (X, G, μ) and (Y, H, ν) are said to be *isomorphic* if there exist a group isomorphism $\xi : G \to H$ and a measure preserving homeomorphism $F : X \to Y$ such that $F \circ g = \xi(g) \circ F$ for all $g \in G$.

Theorem 19. Let $\mathcal{S} = \{S_1, \ldots, S_k\}$ be a set of strong recognizable substitutions. Then the ergodic dynamical systems $(X_{\bar{a}}, T, \mu_{\bar{a}})$ and $(X_{\sigma(\bar{a})}, T, \mu_{\sigma(\bar{a})})$ are isomorphic.

Proof. First note that $S_{a_1} \circ \vec{t} = \lambda_{a_1}\vec{t} \circ S_{a_1}$ and $\xi : T \to T$ defined by $\xi(\vec{t}) = \lambda_{a_1}\vec{t}$ is an isomorphism. By recognizability, S_{a_1} is a bijection from $X_{\sigma(\bar{a})}$ onto $X_{\bar{a}}$. On the other hand, if $x, y \in X_{\sigma(\bar{a})}$ and $d(x, y) < \epsilon$, then $d(S_{a_1}(x), S_{a_1}(y)) < \lambda_{a_1}\epsilon$; similarly, if $x, y \in X_{\bar{a}}$ and $d(x, y) < \epsilon$, then $d(S^{-1}_{a_1}(x), S^{-1}_{a_1}(y)) < \lambda_{a_1}\epsilon$. Hence $S_{a_1} : X_{\sigma(\bar{a})} \to X_{\bar{a}}$ is a homeomorphism. To prove that S_{a_1} is measure preserving is sufficient to prove it for cylinders $X_{P,U}$ with U sufficiently small. By strong recognizability, it is clear that $L_{S_{a_1}(P)}(S^n_{\bar{a}}(D_j)) = L_P(S^{n-1}_{\sigma(\bar{a})}(D_j))$. Then

$$\begin{aligned}\mu_{\bar{a}}\big(S_{a_1}(X^{\sigma(\bar{a})}_{P,U})\big) &= \mu_{\bar{a}}\big(X^{\bar{a}}_{S_{a_1}(P),\lambda_{a_1}U}\big) = \mathrm{freq}_{\bar{a}}\big(S_{a_1}(P)\big)\mathrm{vol}(\lambda_{a_1}U)\\ &= \frac{1}{\lambda^d_{a_1}}\mathrm{freq}_{\sigma(\bar{a})}(P)\lambda^d_{a_1}\mathrm{vol}(U) = \mu_{\sigma(\bar{a})}\big(X^{\sigma(\bar{a})}_{P,U}\big).\end{aligned}$$

□

5.2. **Statistical stability.** The inequality (3) says, in particular, that $\bar{a} \mapsto c_{\bar{a}}(D)$ defines a continuous map $\Sigma \to \mathbb{R}$ for each tile D. In this subsection we extend this result to arbitrary patches x'. As a consequence, we will see that the unique measures $\mu_{\bar{a}}$, although defined in different spaces, also satisfy a certain kind of continuity with respect to $\bar{a} \in \Sigma$.

Recall that the *upper Minkowski dimension* of a subset $H \subset \mathbb{R}^d$ can be defined as

$$\mathcal{D}_H := \inf\{\beta : \operatorname{vol}(H^{+r}) = O(r^{d-\beta}) \text{ as } r \to 0^+\}.$$

Set $d-1 \le \mathcal{D} := \max_i\{\mathcal{D}_{\partial D_i}\} < d$.

Theorem 20. Given $N > 0$, there is $C_N > 0$ such that, for any $\delta > 0$ and any $\bar{a}, \bar{b} \in \Sigma$ with $d_\Sigma(\bar{a}, \bar{b}) < \delta$, we have, for all $x' \in \mathcal{P}^N(X)$,

$$|c_{\bar{a}}(x') - c_{\bar{b}}(x')| < C_N \theta_0^{1/\delta},$$

where $\theta_0 = \theta^{\frac{\mathcal{D}-d}{\mathcal{D}-d+\log_\omega \theta}} < 1$ and θ is given by lemma 14.

Proof. Set $\omega = \min_i\{\omega_i\}$, $t = \max_j\{\operatorname{diam}(D_j)\}$ and $v = \min_j\{\operatorname{vol}(D_j)\}$. We have

$$\frac{\operatorname{vol}\big((\partial S^m_{\bar{a}}(D_i))^{+t}\big)}{\operatorname{vol}(S^m_{\bar{a}}(D_i))} \le \frac{\operatorname{vol}\big((\partial D_i)^{+\frac{t}{\omega^m}}\big)}{\operatorname{vol}(D_i)} = O\Big(1/\omega^{m(d-\mathcal{D})}\Big)$$

as $m \to \infty$, for all $\bar{a} \in \Sigma$. Hence, for some constants C_1 and m_1,

$$N^{\bar{a}}_{x'}(n,m,i,j) < \frac{N \operatorname{vol}((\partial S^m_{\bar{a}}(D_i))^{+t})\big(A^{n-m}_{\sigma^m(\bar{a})}\big)_{ij}}{v} < \frac{C_1 N}{v} \frac{\operatorname{vol}(S^m_{\bar{a}}(D_i))\big(A^{n-m}_{\sigma^m(\bar{a})}\big)_{ij}}{\omega^{m(d-\mathcal{D})}}$$

for all $\bar{a} \in \Sigma$ and $n > m > m_1$. On the other hand,

$$\sum_{i=1}^{l} \frac{\operatorname{vol}(S^m_{\bar{a}}(D_i))\big(A^{n-m}_{\sigma^m(\bar{a})}\big)_{ij}}{\operatorname{vol}(S^n_{\bar{a}}(D_j))} = 1$$

for all j. Hence

$$\sum_{i=1}^{l} \frac{L_{x'}(S^m_{\bar{a}}(D_i))\big(A^{n-m}_{\sigma^m(\bar{a})}\big)_{ij}}{\omega^m_{\bar{a}} \operatorname{vol}(S^{n-m}_{\sigma^m(\bar{a})}(D_j))} \le \frac{L_{x'}(S^n_{\bar{a}}(D_j))}{\operatorname{vol}(S^n_{\bar{a}}(D_j)))} \le \sum_{i=1}^{l} \frac{L_{x'}(S^m_{\bar{a}}(D_i))\big(A^{n-m}_{\sigma^m(\bar{a})}\big)_{ij}}{\omega^m_{\bar{a}} \operatorname{vol}(S^{n-m}_{\sigma^m(\bar{a})}(D_j))} + \frac{C_1 N}{v\omega^{m(d-\mathcal{D})}}.$$

Taking the limit $n \to \infty$ we obtain

$$\sum_{i=1}^{l} \frac{L_{x'}(S^m_{\bar{a}}(D_i))}{\omega^m_{\bar{a}}} c_{\sigma^m(\bar{a})}(D_i) \le c_{\bar{a}}(x') \le \sum_{i=1}^{l} \frac{L_{x'}(S^m_{\bar{a}}(D_i))}{\omega^m_{\bar{a}}} c_{\sigma^m(\bar{a})}(D_i) + \frac{C_1 N}{v\omega^{m(d-\mathcal{D})}}, \tag{12}$$

for all $m > m_1$ and $\bar{a} \in \Sigma$. Set

$$\gamma := 1 - \frac{d-\mathcal{D}}{\log_\omega \theta} > 1, \tag{13}$$

with $\theta < 1$ given by lemma 14. Observe also that

$$L_{x'}(S^m_{\bar{a}}(D_i)) \le \frac{\operatorname{vol}(S^m_{\bar{a}}(D_i))}{\operatorname{vol}(x')} \le \frac{\operatorname{vol}(S^m_{\bar{a}}(D_i))}{Nv} \tag{14}$$

In view of (12), (13), (14) and lemma 14, there is a constant C such that, if, for some $m > 0$, $d_\Sigma(\bar{a}, \bar{b}) < \frac{1}{\gamma m} = \delta$,

$$\begin{aligned}|c_{\bar{a}}(x') - c_{\bar{b}}(x')| &\leq \sum_{i=1}^{l} \frac{L_{x'}(S^m_{\bar{a}}(D_i))}{\omega^m_{\bar{a}}} |c_{\sigma^m(\bar{a})}(D_i) - c_{\sigma^m(\bar{b})}(D_i)| + \frac{C_1 N}{v\omega^{m(d-\mathcal{D})}} \\ &\leq \sum_{i=1}^{l} \frac{\mathrm{vol}(D_i)}{Nv} C'\theta^{(\gamma-1)m} + \frac{C_1 N}{v\omega^{m(d-\mathcal{D})}} \\ &\leq C_N \big(\theta^{\frac{\mathcal{D}-d}{\gamma \log_\omega \theta}}\big)^{1/\delta}\end{aligned}$$

for all $x' \in \mathcal{P}^N(X)$. □

From theorem 13 and theorem 20 we have:

Corollary 21. Given $N > 0$, there is $C_N > 0$ such that, for any $\delta > 0$ and any $\bar{a}, \bar{b} \in \Sigma$ with $d_\Sigma(\bar{a}, \bar{b}) < \delta$, we have, for all $x' \in \mathcal{P}^N(X)$ and all sufficiently small Borel subset $U \subset \mathbb{R}^d$,

$$|\mu_{\bar{a}}(X^{\bar{a}}_{x',U}) - \mu_{\bar{b}}(X^{\bar{b}}_{x',U})| < C_N \theta_0^{1/\delta},$$

where θ_0 is given by theorem 20.

To finalize, we extend the previous corollary:

Theorem 22. Take finite patches $P, Q \in \mathcal{P}(X)$ and measurable sets $U, V \subset \mathbb{R}^d$. There exists $C > 0$ such that, for any $\delta > 0$ sufficiently small and any $\bar{a}, \bar{b} \in \Sigma$ with $d_\Sigma(\bar{a}, \bar{b}) < \delta$, we have

$$|\mu_{\bar{a}}(X^{\bar{a}}_{P,U} \cap X^{\bar{a}}_{Q,V}) - \mu_{\bar{b}}(X^{\bar{b}}_{P,U} \cap X^{\bar{b}}_{Q,V})| < C\theta_0^{\frac{1}{\delta}},$$

where θ_0 is given by theorem 20.

Proof. Let $B(P,U)$ and $B(Q,V)$ be two open balls such that $\mathrm{supp}(\vec{u} + P) \subset B(P,U)$ and $\mathrm{supp}(Q + \vec{v}) \subseteq B(Q,V)$ for all $\vec{u} \in U$ and $\vec{v} \in V$. Consider the set $\mathcal{P}(P, Q, \bar{a})$ of finite patches x' of $X_{\bar{a}}$ such that $B := B(P,U) \cup B(Q,V)$ is contained in the interior of $\mathrm{supp}(x')$ and all tiles of x' intersect B. There are finitely many equivalence classes $\{[x'_1], \ldots, [x'_k]\}$ of such patches, with $x'_i \in \mathcal{P}(P, Q, \bar{a})$. For each $i \in I := \{1, \ldots, k\}$, set

$$U_i = \{\vec{w} : \ \vec{w} + x'_i \in \mathcal{P}(P, Q, \bar{a})\},$$

which is an open subset of $\mathbb{R}^d$. Then we have a disjoint cylinder decomposition

$$X_{\bar{a}} = \bigcup_{i \in I} X^{\bar{a}}_{x'_i, U_i}.$$

The diameter of U_i is smaller then $\max_j\{\mathrm{diam}(D_j)\}$. Take $N_0, \psi > 0$ such that, for all $\bar{a}$ and i, the following hold: $\mathrm{vol}(U_i) \leq \psi$; if $x'_i \in \mathcal{P}^N(X)$ then $N \leq N_0$. For simplicity of exposition we assume that $\mathrm{diam}(U_i) < \eta$ where η is such that any tile contains a ball of radius η. Otherwise, we could decompose each U_i into a fixed number of disjoint subsets satisfying this property.

Consider the subset $W_i \subset U_i$ of vectors $\vec{w}$ such that $\vec{u} + P$ and $\vec{v} + Q$ are contained in $\vec{w} + x'_i$ for some $\vec{u} \in U$ and $\vec{v} \in V$. We have

$$X^{\bar{a}}_{P,U} \cap X^{\bar{a}}_{Q,V} \cap X^{\bar{a}}_{x'_i, U_i} = X^{\bar{a}}_{x'_i, W_i}.$$

Then

$$\mu_{\bar{a}}(X^{\bar{a}}_{P,U} \cap X^{\bar{a}}_{Q,V}) = \sum_{i\in I} \mu_{\bar{a}}(X^{\bar{a}}_{x'_i,W_i}). \tag{15}$$

Take $\bar{b} \in \Sigma$ with $d_\Sigma(\bar{a},\bar{b}) < \delta$. We have a finite disjoint cylinder decomposition

$$X^{\bar{b}}_{P,U} \cap X^{\bar{b}}_{Q,V} = \bigcup_{i\in I, j\in \hat{I}} X^{\bar{b}}_{x'_i,W_i} \cup X^{\bar{b}}_{\hat{x}'_j,\hat{W}_j},$$

where, for each $i \in \hat{I}$, $\hat{x}'_i$ is a patch in $\mathcal{P}(P,Q,\bar{b})$ but not in $\mathcal{P}(P,Q,\bar{a})$. Hence

$$\mu_{\bar{b}}(X^{\bar{b}}_{P,U} \cap X^{\bar{b}}_{Q,V}) = \sum_{i\in I} \mu_{\bar{b}}(X^{\bar{b}}_{x'_i,W_i}) + \sum_{i\in \hat{I}} \mu_{\bar{b}}(X^{\bar{b}}_{\hat{x}'_i,\hat{W}_i}). \tag{16}$$

Equations (15) and (16) give

$$\begin{aligned} |\mu_{\bar{b}}(X^{\bar{b}}_{P,U} \cap X^{\bar{b}}_{Q,V}) - \mu_{\bar{a}}(X^{\bar{a}}_{P,U} \cap X^{\bar{a}}_{Q,V})| & \\ \leq \sum_{i\in I} |\mu_{\bar{b}}(X^{\bar{b}}_{x'_i,W_i}) - \mu_{\bar{a}}(X^{\bar{a}}_{x'_i,W_i})| + \sum_{i\in \hat{I}} \mu_{\bar{b}}(X^{\bar{b}}_{\hat{x}'_i,\hat{W}_i}). & \end{aligned} \tag{17}$$

But, taking account theorem 20, we have

$$\begin{aligned} \sum_{i\in I} |\mu_{\bar{b}}(X^{\bar{b}}_{x'_i,W_i}) - \mu_{\bar{a}}(X^{\bar{a}}_{x'_i,W_i})| &= \sum_{i\in I} |c_{\bar{b}}(x'_i) - c_{\bar{a}}(x'_i)| \mathrm{vol}(W_i) \\ &\leq C|I|\psi\theta_0^{\frac{1}{\delta}}, \end{aligned} \tag{18}$$

for some constant C. On the other hand, it follows from

$$\sum_{i\in I} \mu_{\bar{a}}(X^{\bar{a}}_{x'_i,U_i}) = \sum_{i\in I} \mu_{\bar{b}}(X^{\bar{b}}_{x'_i,U_i}) + \sum_{i\in \hat{I}} \mu_{\bar{b}}(X^{\bar{b}}_{\hat{x}'_i,\hat{U}_i})$$

that

$$\sum_{i\in \hat{I}} \mu_{\bar{b}}(X^{\bar{b}}_{\hat{x}'_i,\hat{U}_i}) \leq \sum_{i\in I} |\mu_{\bar{b}}(X^{\bar{b}}_{x'_i,U_i}) - \mu_{\bar{a}}(X^{\bar{a}}_{x'_i,U_i})| \leq C|I|\psi\theta_0^{\frac{1}{\delta}}. \tag{19}$$

Finally, the result follows from (17), (18) and (19). □

References

[1] F. Durand, Linearly recurrent subshifts have a finite number of non-periodic subshift factors. Ergodic Theory Dynam. Systems, 20, **4** (2000) 1061–1078.

[2] S. Ferenczi, Rank and symbolic complexity. Ergodic Theory Dynam. Systems, 16, **4** (1996) 663–682.

[3] N. P. Frank , A primer on substitution tilings of the Euclidean plane. Expositiones Mathematicae, volume 26, **4** (2008) 295–326.

[4] N. P. Frank and L. Sadun, Fusion: a general framework for hierarchical tilings of $\mathbb{R}^d$. http://arxiv.org/abs/1101.4930.

[5] F. Gähler and G. Maloney, Cohomology of one-dimensional mixed substitution tiling spaces. arXiv:1112.1475 (2011).

[6] C.P.M. Geerse and A. Hof, Lattice gas models on self-similar aperiodic tilings. Rev. Math. Phys. **3** (1991) 163–221.

[7] W.H. Gottschalk, Orbit-closure decomposition and almost periodic properties. Bull. Amer. Math. Soc., **50** (1944) 915–919.

[8] B. Grünbaum and G. C. Shephard, *Tilings and Patterns*. Freeman, New York 1986.

[9] J.-Y. Lee, R. V. Moody, and B. Solomyak, Pure Point Dynamical and Diffraction Spectra, Ann. Henri Poincaré **3** (2002), 1003–1018.
[10] R. Pacheco and H. Vilarinho, Metrics on Tiling Spaces, Local Isomorphism and an Application of Brown's Lemma. arXiv:1202.4902v1 (2012).
[11] C. Radin and M. Wolff, Space Tilings and Local Isomorphism, Geometriae Dedicata **42** (1992), 355–360.
[12] E. A. Robinson, Jr., Symbolic dynamics and tilings of $\mathbb{R}^d$. *Symbolic dynamics and its applications*, volume 60 of Proc. Sympos. Appl. Math., Amer. Math. Soc., Providence 2004, 81–119.
[13] D. Ruelle, *Statistical mechanics: Rigorous results*, W. A. Benjamin, Inc., New York - Amsterdam, 1969.
[14] B. Solomyak, Dynamics of Self-Similar Tilings, Ergodic Theory and Dynamical Systems **17** (1997), 695–738. Errata, Ergodic Theory and Dynamical Systems **19** (1999), 1685.

Universidade da Beira Interior, Rua Marquês d'Ávila e Bolama, 6200-001 Covilhã, Portugal
E-mail address: rpacheco@ubi.pt
URL: http://www.mat.ubi.pt/∼rpacheco

Universidade da Beira Interior, Rua Marquês d'Ávila e Bolama, 6200-001 Covilhã, Portugal
E-mail address: helder@ubi.pt
URL: http://www.mat.ubi.pt/∼helder